



MEF2C-related syndrome





This guide is not meant to take the place of medical advice.

Please consult with your doctor about your genetic results and health care choices. The information in this guide was up to date at the time it was written in 2020. But new information may come to light with new research. You may find it helpful to share this guide with friends and family members or doctors and teachers of the person who has MEF2C-related syndrome.





What is MEF2C-related syndrome?

MEF2C-related syndrome happens when there are changes to the MEF2C gene. These changes can keep the gene from working as it should.

Key role

The MEF2C gene is important for muscle and brain cells.

Symptoms


Because the MEF2C gene is important in the development and function of brain cells, many people who have MEF2C-related syndrome have:

- Intellectual disability
- Speech difficulties or no speech
- Delayed walking
- Autism or features of autism

What causes MEF2C-related syndrome?

Our genes contain the instructions, or code, that tell our cells how to grow, develop, and work. Every child gets two copies of the MEF2C gene: one copy from their mother, from the egg, and one copy from their father, from the sperm. In most cases, parents pass on exact copies of the gene to their child. But the process of copying genes is not perfect. A change in the genetic code can lead to physical issues, developmental issues, or both.

Sometimes a random change happens in the sperm or egg. This change to the genetic code is called a 'de novo', or new, change. The child can be the first in the family to have the gene change.




De novo changes can take place in any gene. We all have some de novo changes, most of which don't affect our health. But because MEF2C plays a key role in development, de novo changes in this gene can have a meaningful effect.

Research shows that MEF2C-related syndrome is often the result of a de novo change in MEF2C. Many parents who have had their genes tested do not have the MEF2C gene change found in their child who has the syndrome. In some cases, MEF2C-related syndrome happens because the gene change was passed down from a parent. This is called dominant inheritance.



Dominant inheritance
Children have a 50% chance of inheriting the genetic change



Why does my child or I have a change in the MEF2C gene?

No parent causes their child's MEF2C-related syndrome. We know this because no parent has any control over the gene changes that they do or do not pass on to their children. Please keep in mind that nothing a parent does before or during the pregnancy causes this to happen. The gene change takes place on its own and cannot be predicted or stopped.





What are the chances that other family members or future children will have MEF2C-related syndrome?

Each family is different. A geneticist or genetic counselor can give you advice on the chance that this will happen again in your family.

The risk of having another child who has MEF2C-related syndrome depends on the genes of both birth parents.

- If neither birth parent has the same gene change found in their child, the chance of having another child who has the syndrome is on average 1 percent. This 1 percent chance is higher than the chance of the general population. The increase in risk is due to the very unlikely chance that more of the mother's egg cells or the father's sperm cells carry the same change in the gene.
- If one birth parent has the same gene change found in their child, the chance of having another child who has the syndrome is 50 percent.

For a symptom-free sibling, a brother or sister, of someone who has MEF2C-related syndrome, the risk of having a child who has the syndrome depends on the symptom-free sibling's genes and their parents' genes.

- If neither parent has the same gene change found in their child who has the syndrome, the symptom-free sibling has a nearly 0 percent chance of having a child who has MEF2C-related syndrome.
- If one birth parent has the same gene change found in their child who has the syndrome, the symptom-free sibling has a small chance of also having the same gene change. If the symptom-free sibling has the same gene change as their sibling who has the syndrome, the symptom-free sibling's chance of having a child who has MEF2C-related syndrome is 50 percent.

For a person who has MEF2C-related syndrome, the risk of having a child who has the syndrome is about 50 percent.


How many people have MEF2C-related syndrome?

As of 2020, about 30 people in the world with changes in the MEF2C gene had been described in medical research. The first case of MEF2C-related syndrome was described in 2010. Scientists expect to find more people who have the syndrome as access to genetic testing improves.

Do people who have MEF2C-related syndrome look different?

People who have MEF2C-related syndrome may look different. Appearance can vary and can include some but not all of these features:

- **Broad forehead**
- **Prominent ear lobes**
- **Tented upper lip**





How is MEF2C-related syndrome treated?

Scientists and doctors have only just begun to study MEF2C-related syndrome. At this point, there are no medicines designed to treat the syndrome. A genetic diagnosis can help people decide on the best way to track the condition and manage therapies. Doctors can refer people to specialists for:

- Physical exams and brain studies.
- Genetics consults.
- Development and behavior studies.
- Other issues, as needed.

A developmental pediatrician, neurologist, or psychologist can follow progress over time and can help:

- Suggest the right therapies. This can include physical, occupational, speech, or behavioral therapy.
- Guide individualized education plans (IEPs).

Specialists advise that therapies for MEF2C-related syndrome should begin as early as possible, ideally before a child begins school.

If seizures happen, consult a neurologist. There are many types of seizures, and not all types are easy to spot. To learn more, you can refer to resources such as the Epilepsy Foundation's website: www.epilepsy.com/learn/types-seizures.

MEF2C-related syndrome is very rare. Doctors and scientists have just recently begun to study it. As of 2020, studies had described fewer than 30 people who have MEF2C-related syndrome.

This section includes a summary of information from major published articles. It highlights how many people have different symptoms. To learn more about the articles, see the [Sources and references](#) section of this guide.

Behavior and development concerns linked to MEF2C-related syndrome

Learning

Most people have **intellectual disability**, which is often severe.



Speech

Many people have **language difficulties**.

Behavior

Some people show **features of autism**, such as **poor eye contact**.



Medical and physical concerns linked to MEF2C-related syndrome

Brain

Seizures during infancy or childhood are common. This can include different types of seizures.




Sitting and walking

Many people have **motor delays**. Most people eventually learn to walk, ranging in age from 22 months to 8 years.

Muscle tone

Some people have **low muscle tone**.



Feeding and digestion issues

Some people have **feeding difficulties**.



Where can I find support and resources?

Simons Searchlight is another research program sponsored and run by the Simons Foundation Autism Research Initiative, also known as SFARI. As part of the next step in your research journey, Simons Searchlight offers you the opportunity to partner with scientists and other families who have the same gene change. Simons Searchlight is a registry for more than 200 genetic changes that are associated with neurodevelopmental conditions, including autism spectrum disorder. Simons Searchlight makes it easier for researchers to access the information they need to advance research on a condition.

To register for Simons Searchlight, go to the Simons Searchlight website at www.simonssearchlight.org and click “Join Us Today”.

- **Learn more about Simons Searchlight**
www.simonssearchlight.org/frequently-asked-questions
- **Simons Searchlight webpage with more information on MEF2C**
www.simonssearchlight.org/research/what-we-study/mef2c
- **Simons Searchlight MEF2C Facebook community**
www.facebook.com/groups/1027885070966893

Sources and References

The content in this guide comes from published studies about MEF2C-related syndrome. Below you can find details about each study, as well as links to summaries or, in some cases, the full article.

- Rocha H. *et al.* *European Journal of Medical Genetics*, **59**, 478-482, (2016). MEF2C haploinsufficiency syndrome: report of a new MEF2C mutation and review
www.ncbi.nlm.nih.gov/27255693
- Borlot F. *et al.* *Seizure*, **67**, 86-90, (2019). MEF2C-related epilepsy: delineating the phenotypic spectrum from a novel mutation and literature review
[www.seizure-journal.com/article/S1059-1311\(19\)30005-6/fulltext](http://www.seizure-journal.com/article/S1059-1311(19)30005-6/fulltext)



© Simons Foundation
Last updated 2020

